

# Coulomb blockade in metallic grains at large conductance: beyond the zero-dimensional limit

I.S. Beloborodov and A.V. Andreev

*Bell Laboratories, Lucent Technologies, Murray Hill, NJ 07974*

*Department of Physics, University of Colorado, CB 390, Boulder, CO 80390*

(February 1, 2008)

We consider mesoscopic fluctuations of the Coulomb blockade oscillations in a disordered metallic grain in an external magnetic field coupled to a metallic lead by a tunneling contact with a large conductance  $g_T$  and capacitively coupled to a gate. For a finite mean level spacing  $\delta$  in the grain and at temperatures  $T \gg \delta$  we obtain the oscillatory part of the correlator of thermodynamic potentials at different values of the gate voltage. The correlations decay algebraically with the gate voltage difference in contrast to the exponential decay that would result from the mean-field treatment of the Coulomb interaction. The results are valid beyond the zero-dimensional limit and describe the crossover between the unitary and orthogonal ensembles.

PACS numbers: 73.23Hk, 73.40 Gk, 73.21.La

Coulomb blockade is one of the most striking manifestations of electron interactions at low temperatures [1]. It can be observed by measuring, say the charge of a small metal particle connected by a tunneling contact to a metallic lead and capacitively coupled to a metallic gate which is maintained at voltage  $V_g$ , as in Fig. 1.

In the weak tunneling regime the charge in the grain is nearly quantized at low temperatures and exhibits a characteristic step-like dependence on the gate voltage  $V_g$ . This leads to the Coulomb blockade oscillations in the gate voltage dependence of various physical quantities. The amplitude of these oscillations decreases with increasing transparency of the contact and depends not only on the total conductance of the contact but on the individual transparencies of the tunneling channels.

In the strong tunneling regime most of the studies of the Coulomb blockade concentrated on the limit of large grains with the vanishing mean level spacing  $\delta$ . In this limit the Coulomb blockade oscillations of the thermodynamic quantities vanish at perfect transmission [2,3]. In the case of a multi-channel tunneling contact the Coulomb blockade oscillations at large conductance have been shown to be exponentially small in the dimensionless tunneling conductance  $g_T$  [4,5]. The case of a diffusive random contact was considered in Refs. [6,7]. At  $\delta = 0$  the mesoscopic fluctuations of the Coulomb blockade oscillations due to disorder in the grain vanish.

The case of strong tunneling and finite  $\delta$  was studied in Ref. [8] where it was shown that the Coulomb blockade oscillations remain finite even at perfect transmission and exhibit strong mesoscopic fluctuations due to the presence of disorder in the grain. This study utilized the bosonization method of Ref. [3] which is applicable to contacts with weakly reflecting channels.

Recently, much interest was focussed on granulated disordered metals [9,10], which can be viewed as extended networks of metallic grains in the strong tunneling

regime. The low temperature regime, where mesoscopic fluctuations at finite  $\delta$  become observable is well within experimental reach [10]. It is therefore desirable to generalize the study of charging effects at finite  $\delta$  and strong tunneling [8] to granulated metals. The formalism of Refs. [3,8] is well suited to treat individual grains but is difficult to generalize to extended disordered systems. On the other hand such systems are conveniently described within the  $\sigma$ -model formalism. It is therefore important to incorporate the non-perturbative Coulomb blockade effects into the framework of the  $\sigma$ -model approach.




FIG. 1. Schematic drawing of a disordered metallic grain coupled by a tunneling contact to a metallic lead. The charge of the grain is controlled by the gate voltage  $V_g \sim q$

In this Letter we show that mesoscopic fluctuations of the Coulomb blockade oscillations in a metallic grain with finite  $\delta$  in the strong tunneling regime can be studied using the  $\sigma$ -model. We concentrate on the thermodynamic quantities of a grain of size  $L$  with the Thouless energy  $E_T = D/L^2$ , where  $D$  is the diffusion constant, placed in an external magnetic field  $H$ . The grain is connected to a metallic lead by a non-random multi-channel tunneling contact with the dimensionless conductance  $g_T = 2\pi\hbar/e^2 R \gg 1$  where  $R$  is the resistance of the contact. It is also capacitively coupled to a gate, as in Fig. 1. We assume that the electron-electron inter-

action in the lead may be neglected due to screening and treat the Coulomb interaction of electrons in the grain within the framework of the constant interaction model,

$$\hat{H}_C = E_C (\hat{N} - q)^2. \quad (1)$$

Here  $E_C$  is the charging energy,  $\hat{N}$  is the operator of the number of electrons in the grain, and  $q \propto V_g$  is the number of electrons that minimizes the electrostatic energy of the grain. We consider the case  $E_T < E_C$ , and describe the crossover between the unitary and orthogonal ensembles. The case  $E_T > E_C$  for the unitary ensemble was recently studied in Ref. [11] by a different method.

The single particle levels in the grain are broadened due to tunneling into the lead. We assume that tunneling contact is broad, so that each state  $x$  in the dot is coupled to many lead states  $a$ . In this case the probability distribution  $P(\Gamma_x)$  of level half-widths  $\Gamma_x$  is sharply peaked about the mean value  $\Gamma$ . We can therefore neglect the fluctuations of  $P(\Gamma_x)$  and consider them to be equal to the mean value  $\Gamma$  which can be expressed through the dimensionless conductance of the contact  $g_T$  as  $\Gamma = \frac{g_T \delta}{8\pi}$ . The temperature is assumed to satisfy the conditions  $\delta \ll T \ll E_C$ .

The main results of this paper are the following expressions for the oscillatory parts of the *ensemble averaged* thermodynamic potential  $\langle \Omega(q) \rangle_{\text{osc}}$  and of the irreducible correlator  $\langle \langle \Omega(q) \Omega(q') \rangle \rangle_{\text{osc}} = \langle \Omega(q) \Omega(q') \rangle_{\text{osc}} - \langle \Omega(q) \rangle_{\text{osc}} \langle \Omega(q') \rangle_{\text{osc}}$  at  $\delta |q - q'| \gg \Gamma + T$ ,

$$\langle \Omega(q) \rangle_{\text{osc}} = -\tilde{E}_C \ln \left[ \frac{E_C}{T + \Gamma} \right] \cos(2\pi q), \quad (2a)$$

$$\langle \langle \Omega(q) \Omega(q') \rangle \rangle_{\text{osc}} = \frac{1}{2} \tilde{E}_C^2 \Re \sum_{\rho=0,1} \left[ e^{2\pi i(q-q')} B_\rho \right]. \quad (2b)$$

Here  $\tilde{E}_C = \frac{g_T E_C}{\pi^2} e^{-g_T/2}$  and  $B_\rho$  can be conveniently expressed through the quantity  $a_\rho = i\delta(q - q')/2 + \rho/\tau_H$ , with the cooperon gap  $1/\tau_H = DeH/\hbar c$ ,

$$B_\rho = \begin{cases} \left( \frac{4\Gamma}{a_\rho} \right)^2 \ln^2 \frac{a_\rho}{\Gamma}, & |a_\rho| \ll E_T, \\ \frac{4^{3-d}\Gamma^2}{\pi E_T^{d/2} a_\rho^{2-d/2}} \ln^2 \frac{a_\rho}{\Gamma}, & |a_\rho| \gg E_T. \end{cases} \quad (2c)$$

Here the index  $\rho$  labels the diffuson,  $\rho = 0$ , and the cooperon,  $\rho = 1$ , degrees of freedom and  $d = 2, 3$  is the dimensionality of the grain.

The results (2) were obtained with logarithmic accuracy in the one instanton approximation and are valid for  $T \gg g_T^2 E_C \ln(E_C/\Gamma) \exp(-g_T/2)$ . The Zeeman splitting of energy levels was neglected. Equation (2a) coincides with the result for the zero-dimensional grain [11]. At  $q - q' \rightarrow \infty$  the function  $B_\rho$  in Eq. (2c) tends to zero, and the correlator  $\langle \Omega(q) \Omega(q') \rangle_{\text{osc}}$  factorizes;  $\langle \Omega(q) \Omega(q') \rangle_{\text{osc}} = \langle \Omega(q) \rangle_{\text{osc}} \langle \Omega(q') \rangle_{\text{osc}}$ .

Below we outline the derivation of Eq. (2). We consider an ensemble of disordered metallic grains in which

electrons move in the presence of a random impurity potential  $V(\mathbf{r})$  which is taken to be white noise random potential with the variance  $\langle V(\mathbf{r}) V(\mathbf{r}') \rangle = \delta(\mathbf{r} - \mathbf{r}')/2\pi\nu\tau$ , where  $\nu$  is the density of states. The correlator of the thermodynamic potentials at different values of the gate voltage  $q$  and  $q'$  can be written as

$$\langle \Omega(q) \Omega(q') \rangle = T^2 \frac{\partial^2}{\partial \alpha \partial \alpha'} \langle Z^\alpha(q) Z^{\alpha'}(q') \rangle |_{\alpha, \alpha' \rightarrow 0}, \quad (3)$$

Here  $\langle Z^\alpha(q) Z^{\alpha'}(q') \rangle$  is the ensemble averaged product of the replicated partition functions at different values of the gate voltage. For  $\delta \ll T \ll E_C$  it can be expressed as a functional integral over the  $\hat{Q}$ -matrices, arising from disorder averaging, and over the phases  $\phi_j$  which arise from decoupling the Coulomb interaction (1) via the Hubbard-Stratonovich transformation, as follows:

$$\langle Z^\alpha(q) Z^{\alpha'}(q') \rangle = \sum_{\{W\}} \langle Z^\alpha Z^{\alpha'} \rangle_W e^{-2\pi i \sum_{j=1}^{\alpha+\alpha'} q_j W_j}, \quad (4a)$$

$$\langle Z^\alpha Z^{\alpha'} \rangle_W = \int d\hat{Q} d[\phi] e^{-F[\hat{Q}, \phi] - \sum_{j=1}^{\alpha+\alpha'} \left[ \frac{E_C q_j^2}{T} + \int_0^\beta \frac{\dot{\phi}_j^2 d\tau}{4E_C} \right]}, \quad (4b)$$

$$F[\hat{Q}, \phi] = \text{Tr} \left[ \frac{\pi\nu}{4\tau} \hat{Q}^2 - \ln \left( i\hat{J} + \frac{\vec{\partial}^2}{2m} + \frac{i\hat{Q}}{2\tau} \right) \right]. \quad (4c)$$

Equations (4) were obtained by taking the integral over the static component of the field  $\phi$  by the saddle point approximation which is valid for  $T \gg \delta$  [12]. The resulting expression, Eq. (4a), is written as a sum over the sets of winding numbers  $\{W\}$  [16] which label distinct topological classes of the phase,  $\phi_j(\beta) = \phi_j(0) + 2\pi W_j$ . The treatment of the cooperon degrees of freedom requires doubling the number of fermion fields. The covariant derivative  $\vec{\partial}$  in Eq. (4c) is expressed through the time-reversal symmetry breaking matrix  $\tau_3$  [14] as  $\vec{\partial} = \nabla + ie\mathbf{A}\hat{\tau}_3/c$ , where  $\mathbf{A}$  is the vector potential, and  $\text{Tr}$  denotes the trace taken over the spin,  $\sigma$ , coordinate,  $r$ , replica,  $i, j$ , and Matsubara,  $n, m$ , as well as the time-reversal indices. The matrix  $\hat{J}$  in Eq. (4c) is defined as

$$\hat{J} \equiv \left[ \hat{\varepsilon} - \frac{iq_j \delta}{2} + \Gamma e^{-i\phi_j(\tau)} \Lambda_{\tau, \tau'} e^{i\phi_j(\tau')} \right] \delta_{ij} \delta_{\sigma, \sigma'}, \quad (5)$$

where  $\hat{\varepsilon} = i\partial_\tau$  is diagonal in the Matsubara frequencies,  $\hat{\varepsilon} = \delta_{nm} \varepsilon_n = \delta_{nm} \pi T(2n+1)$ , and  $\Lambda_{\tau, \tau'} = -i/\sin[\pi T(\tau - \tau')]$ . Finally,  $q_j = q$  for  $1 \leq j \leq \alpha$  and  $q_j = q'$  for  $\alpha < j \leq \alpha + \alpha'$ .

At frequencies smaller than the inverse mean free time  $1/\tau$  the effective low-energy theory may be written as a functional integral over the saddle point manifold of  $\hat{Q}$ -matrices [15]. The low-energy degrees of freedom are usually parameterized as  $\hat{Q} = U \hat{Q}_0 U^{-1}$ , where the matrix  $\hat{Q}_0$  is diagonal in the Matsubara space, and represents

the saddle point of the free energy  $F[\hat{Q}, \phi]$  in Eq. (4c) at  $\hat{\phi} = 0$ . At frequencies below the Fermi energy it is given by  $\hat{Q}_0 = \Lambda$ . For our present purposes, however, it is convenient to write the  $\hat{Q}$ -matrix as  $\hat{Q} = \hat{T}\hat{\lambda}\hat{T}^{-1}$ , where  $\hat{\lambda}$  is the saddle point with the lowest free energy  $F[\hat{Q}, \phi]$  for a given  $\hat{\phi}$  [13]. It is given by the equation

$$\hat{\lambda} = \frac{i}{\pi\nu} \int \frac{d^d p}{(2\pi)^d} \left( i\hat{J} + \frac{(p - \frac{e\mathbf{A}\hat{\tau}_3}{c})^2}{2m} + \hat{\mu}(\hat{q}) + \frac{i\hat{\lambda}}{2\tau} \right)^{-1}. \quad (6)$$

Note that since the operator  $\hat{J}$  is coordinate-independent, the saddle point matrix  $\hat{\lambda}$  is uniform in space. This is a consequence of the constant interaction model, Eq. (1).

At  $\phi_j(\tau) = 2\pi T\tau W_j$  the matrix  $\hat{J}$  is diagonal in the basis of Matsubara frequencies. Since it is Hermitian it may be diagonalized in some different basis for other configurations of  $\phi_j(\tau)$  as well:  $\hat{J} = \delta_{nm}\delta_{ij}\delta_{\sigma,\sigma'} J_n^j$ , where  $n, m$  can be identified with the Matsubara indices only at  $\phi_j(\tau) = 2\pi T\tau W_j$ . The saddle point  $\hat{\lambda}$ , Eq. (6) is also diagonal in this basis and in the low-frequency sector is given by  $\hat{\lambda} = \text{sgn}\varepsilon_n\delta_{nm}\delta_{ij}\delta_{\sigma,\sigma'}$ , where  $\varepsilon_n$  should be understood as the Matsubara frequency corresponding to  $J_n^j$  at  $\phi_j(\tau) = 2\pi T\tau W_j$ .

At  $T \gg \delta$  the integration over the soft modes of  $\hat{Q}$  in Eq. (4b) may be carried out by expressing the rotation matrices  $\hat{T}$  in terms of the generators,  $\hat{T} = \exp(iP)$ , and expanding the free energy  $F[\hat{Q}, \phi]$  to second order in  $P$ :

$$F[\hat{Q}, \phi] = F[\hat{\lambda}, \phi] + \frac{\pi}{2\delta} \sum_{\rho=0,1} \sum_{\sigma,\sigma'} \sum_{i,j;n,m} |P_{ki;\sigma\sigma'}^{mn}(\rho)|^2 \times (D\mathbf{p}^2 + J_n^i - J_m^j + \rho/\tau_H). \quad (7)$$

The prime over the sum in Eq. (7) and below indicates that the summation goes only over such pairs  $n, m$  that  $\varepsilon_n > 0$  and  $\varepsilon_m < 0$ . The Gaussian integration over  $P$  in Eq. (4b) then results in the following expression

$$\Delta_{\{W\}} = \prod_{i,j;\rho} \prod_{\mathbf{p},m,n} \frac{\delta^4}{(J_n^i - J_m^j + D\mathbf{p}^2 + \rho/\tau_H)^4}, \quad (8)$$

where the power 4 arises from taking the product over the spin indices.

The largest term in the sum over  $\{W\}$  in Eq. (4a) has a monotonic dependence on the gate voltages,  $q$  and  $q'$ , and corresponds to  $\{W\} = \{0\}$  with all  $W_j = 0$ . The leading oscillatory contributions to this sum arise from the terms with  $\{W\} = \{l, l'\}$ , with  $l, l' = \pm 1$ , having only one non-zero  $W_j = l$  for  $1 \leq j \leq \alpha$  and only one non-zero  $W_j = l'$  for  $\alpha < j \leq \alpha + \alpha'$ . The dissipative free energy  $F[\hat{\lambda}, \phi]$  in Eq. (7) has a minimum when  $\phi_j = 0$  in the replicas with  $W_j = 0$ . In the replica with  $W_j = l$  the instanton phase configuration [17] which minimizes the free energy may be written in terms of a complex variable  $z$  [6],

$$\exp(i\phi_z) = \frac{u - z}{1 - uz^*}, \quad u = \exp(2\pi iT\tau), \quad (9)$$

whereas for  $W_j = l'$   $\phi_j$  is given by Eq. (9) with  $z \rightarrow z'$ . The free energy  $F[\hat{\lambda}, \phi] = g_T$  on the instanton configuration (9) and is independent of  $z$  and  $z'$  which parameterize the widths and the positions of the instantons in replicas with  $W_j = l, l'$  and satisfy the inequalities:  $|z|, |z'| < 1$  for  $l, l' = 1$ , and  $|z|, |z'| > 1$  for  $l, l' = -1$ . We need to integrate over the fluctuations of  $\phi$  around the instanton configuration Eq. (9). To this end we write  $\phi = \phi_z + \tilde{\phi}$ , where  $\tilde{\phi}$  represents the massive modes. The integration over the zero modes should be performed with the measure  $d^2 z / (1 - |z|^2)$  for  $|z| < 1$ . For  $|z| > 1$  the inversion transformation  $z \rightarrow 1/z$  again maps the integration domain to  $|z| < 1$  with the integration measure  $d^2 z / (1 - |z|^2)$ . All other fluctuations  $\tilde{\phi}$  have a large mass of order  $g_T$  and can be integrated out in the gaussian approximation. Moreover, in carrying out this integration one may replace the determinants  $\Delta[\phi]$  in Eq. (8) by their values  $\Delta[\phi_z]$  on the instanton configurations (9) due to their weak dependence on  $\tilde{\phi}$ . We then obtain

$$\frac{\langle Z^\alpha(q) Z^{\alpha'}(q') \rangle}{\langle Z^\alpha(q) Z^{\alpha'}(q') \rangle_{\{0\}}} = 1 + \alpha\alpha' \sum_{l,l'=\pm 1} e^{2\pi i(lq+l'q')} \times \int_{|z|,|z'|<1} \frac{d^2 z d^2 z' f(z) f(z')}{(1 - |z|^2)(1 - |z'|^2)} \frac{\Delta_{\{l,l'\}}[\phi_z]}{\Delta_{\{0\}}[0]}, \quad (10)$$

where  $f(z)$  is given by the following expression [11]

$$f(z) = \frac{g_T^2 E_C \exp\left\{-\frac{2\pi^2 T |z|^2}{E_C(1 - |z|^2)}\right\}}{2\pi^3 T \left[1 + \frac{\Gamma}{\pi T} (1 - |z|^2)\right]}. \quad (11)$$

With the aid of Eq. (8) the determinant  $\Delta_{\{l,l'\}}[\phi_z]$  in Eq. (10) can be expressed through the eigenvalues of the operator  $\hat{J}$  on the instanton configuration (9). To this end it is convenient to express the operator  $\hat{J}$ , Eq. (5), in terms of the complex variable  $u$  in Eq. (9),

$$\frac{\hat{J} + \frac{iq\delta}{2}}{2\pi T} = -u\partial_u - \frac{i\Gamma}{2\pi^2 T} \oint \frac{du'}{u'} \frac{1 - uz^*}{u - z} \frac{\sqrt{uu'}}{u - u'} \frac{u' - z}{1 - u'z^*}. \quad (12)$$

The integral in this equation is taken over the unit circle,  $|u'| = 1$ , and the operator  $\hat{J}$  is acting in the space of functions spanned by the fermionic Matsubara frequencies  $u^{-n-1/2}$ . In the case of the instantons,  $l, l' = 1$ , (anti-instantons,  $l, l' = -1$ ) the functions  $u^{-n-1/2}$  with negative (positive) Matsubara frequencies, are eigenfunctions of  $\hat{J}$  with eigenvalues  $J_n = \varepsilon_n - \Gamma - iq\delta/2$ . All other eigenvalues  $J_n$  can be found from the following equation [11]

$$\frac{1}{1 - |z|^2} = \sum_{k=0}^{\infty} \frac{2\Gamma|z|^{2k}}{\mp(J_n + iq\delta/2) + \varepsilon_k + \Gamma}, \quad (13)$$

where the top (bottom) sign in the right hand side describes the instanton (anti-instanton) case. For  $\delta|q - q'| \gg \Gamma + T$  the integral in Eq. (10) is dominated by the region of short instantons,  $1 - |z|^2 \ll \frac{T}{\Gamma + T}$  and  $1 - |z'|^2 \ll \frac{T}{\Gamma + T}$ . In this case the solutions of Eq. (13) are given by  $J_n = -i\delta q/2 \pm [\varepsilon_n + \Gamma - 2\Gamma|z|^{2n}(1 - |z|^2)]$ ,  $n \geq 0$ . Next, we use these eigenvalues to evaluate the determinant in Eq. (8). Since the second term in Eq. (10) is multiplied by  $\alpha\alpha'$ , to obtain the correlator in Eq. (3) we only need to evaluate the ratio of the determinants in the second term in the limit  $\alpha, \alpha' \rightarrow 0$ . It is not difficult to show, see appendix C of Ref. [11], that in the limit  $\alpha, \alpha' \rightarrow 0$ ,  $\Delta_{\{0\}}[\phi_z]$ ,  $\Delta_{\{1,1\}}[\phi_z]$  and  $\Delta_{\{-1,-1\}}[\phi]$  are equal to unity, and  $\Delta_{\{1,-1\}}[\phi_z]$  is given by

$$\frac{\Delta_{\{1,-1\}}}{\Delta_{\{0\}}} = e^{\sum_{\rho, \mathbf{p}} \sum_{n, m \geq 0} \frac{16\Gamma^2 |z|^{2n} (1 - |z|^2) |z'|^{2m} (1 - |z'|^2)}{(\varepsilon_n + \varepsilon_m + D\mathbf{p}^2 + a_\rho)^2}}, \quad (14)$$

with  $a_\rho$  defined below Eq. (2b). The ratio  $\frac{\Delta_{\{1,1\}}[\phi_z]}{\Delta_{\{0\}}}$  is given by the complex conjugate of Eq. (14). Using Eqs. (14), (10) and (3) we obtain Eq. (2).

In conclusion we note that: i) Equation (2) was obtained with logarithmic accuracy, and is valid for temperatures  $\delta \ll T \ll E_C$ . To derive it we also neglected the instanton-instanton interactions, which is valid for  $T \gg g_T^2 E_C \ln \frac{E_C}{T} \exp(-gT/2)$ .

ii) The results for the orthogonal and the unitary ensembles are obtained from Eq. (2) by taking the limits  $\tau_H \rightarrow \infty$  and  $\tau_H \rightarrow 0$  respectively. If the Zeeman splitting is taken into account the degeneracy of diffusions/cooperons in Eq. (2c) corresponding to different spin projections of the particles is lifted: two of them remain unchanged, and the other two acquire a correction  $i\delta(q - q') \rightarrow i\delta(q - q') \pm i2\mu H$ , where  $\mu$  is the Bohr magneton.

iii) The correlator of differential capacitances, or thermodynamic densities of states, may be obtained from Eq. (2b) by differentiating it with respect to  $q$  and  $q'$ . The correlations fall off as  $|q - q'|^{\frac{d}{2}-2} \ln^2 |q - q'|$  at  $|q - q'| \gg E_T/\delta$ . Equations (2b) and (2c) demonstrate the inadequacy of the mean-field treatment of the Coulomb interaction for the description of the Coulomb blockade oscillations. Under such treatment the Coulomb interaction in the grain would merely result in the linear  $q$ -dependence of the self-consistent chemical potential for the non-interacting problem. In contrast with Eq. (2) the corresponding oscillatory part of the correlator of densities of states would decay exponentially  $\propto |q - q'|^{-4} \exp \left\{ -\gamma_d \left( \frac{\delta|q - q'|}{E_T} \right)^{d/2} \right\}$  [18], where  $\gamma_d$  is a numerical coefficient.

iv) The mesoscopic fluctuations of the Coulomb blockade oscillations in the strong tunneling regime were first studied in Ref. [8]. The single particle properties of the dot in this approach are described in terms of the scattering matrix at the contact. The long range correlations obtained in Ref. [8] resulted from neglecting the dependence of the scattering matrix on the gate voltage.

v) In the  $\sigma$ -model approach the Coulomb blockade oscillations are described by the instantons of the  $\phi$ -field [17]. Their mesoscopic fluctuations are described by the fluctuations of the  $Q$ -matrix about the instanton configurations. The problem therefore reduces the evaluation of diffusions on the instanton configurations of the  $\phi$ -field.

We thank I. Aleiner, A. Kamenev, A. Larkin, K. Matveev and Yu. Nazarov for helpful discussions. We gratefully acknowledge of the Ruhr-Universität, Bochum, Germany, where part of this work was performed. This research was sponsored by the Grants DMR-9984002, BSF-9800338 and SFB 237, and by the A.P. Sloan and the Packard Foundations.

- 
- [1] L. P. Kouwenhoven *et al.*, *Proceedings of the Advanced Study Institute on Mesoscopic Electron Transport*, Eds. L. Sonh, L. P. Kouwenhoven and G. Schön (Kluwer, Series E, 1997).
  - [2] K. Flensberg, Phys. Rev. B **48**, 11156 (1993).
  - [3] K. A. Matveev, Phys. Rev. B **51**, 1743 (1995).
  - [4] S. V. Panyukov and Zaikin, Phys. Rev. Lett. **67**, 3168 (1991).
  - [5] X. Wang and H. Grabert, Phys. Rev. B **53**, 12 621 (1996).
  - [6] Y. V. Nazarov, Phys. Rev. Lett. **82**, 1245 (1999).
  - [7] A. Kamenev, Phys. Rev. Lett. **85**, 4160 (2000).
  - [8] I. L. Aleiner and L. I. Glazman, Phys. Rev. B **57**, 9608 (1998); I. L. Aleiner, P. W. Brouwer and L. I. Glazman, cond-mat/0103008.
  - [9] I. S. Beloborodov, *et al.*, Phys. Rev. B **63**, 115109 (2001).
  - [10] A. Gerber, *et al.*, Phys. Rev. Lett. **78**, 4277 (1997).
  - [11] I. S. Beloborodov and A. V. Andreev, cond-mat/0103075.
  - [12] A. Kamenev and Y. Gefen, Phys. Rev. B **54**, 5428 (1996).
  - [13] A. Kamenev and A. Andreev, Phys. Rev. B **60**, 2239 (1999).
  - [14] K. B. Efetov, *Supersymmetry in Disorder and Chaos*, Cambridge University Press, New York (1997).
  - [15] A. M. Finkelstein, Sov. Phys. JETP **57**, 97 (1983).
  - [16] V. Ambegaokar, U. Eckern, and G. Schön, Phys. Rev. Lett. **48**, 1745 (1982).
  - [17] S. E. Korshunov, Sov. Phys. JETP Lett., **45**, 434 (1987).
  - [18] A. V. Andreev and B. L. Altshuler, Phys. Rev. Lett. **75**, 902 (1995); A. V. Andreev, B. D. Simons, and B. L. Altshuler, J. Math. Phys. **37**, 4968 (1996); A. Kamenev and M. Mezard, Phys. Rev. B **60**, 3944 (1999).